arXiv:0910.1891v1 [nucl-th] 10 Oct 2009

Longitudinal boost-invariance of charge balance function in hadron-hadron and nucleus-nucleus collisions

Li Na, Li Zhiming, and Wu Yuanfang
Key Laboratory of Quark and Lepton Physics (Huazhong Normal University),Ministry of Education
Institute of Particle Physics, Huazhong Normal University, Wuhan 430079, China

Using Monte Carlo generators of the PYTHIA model for hadron-hadron collisions and a multi-phase transport (AMPT) model for nucleus-nucleus collisions, the longitudinal boost-invariance of charge balance function and its transverse momentum dependence are carefully studied. It shows that the charge balance function is boost-invariant in both p+p and Au+Au collisions in these two models, consistent with experimental data. The balance function properly scaled by the width of the pseudorapidity window is independent of the position or the size of the window and is corresponding to the balance function of the whole pseudorapidity range. This longitudinal property of balance function also holds for particles in small transverse momentum ranges in the PYTHIA and the AMPT default models, but is violated in the AMPT with string melting. The physical origin of the results are discussed.

PACS numbers: 25.75.Gz,25.75.Ld

I. INTRODUCTION

Charge balance function (BF) has been widely used as an effective exploring for the hadronization scheme in hadron-hadron collisions at the ISR energies [1] and $e^+ + e^-$ annihilations at PETRA energies [2]. Recently, the charge BF gains special attentions in clocking hadronization at relativistic heavy-ion collisions. A narrowing of the BF is suggested as a signature for delayed hadronization [3, 4].

The dependence of the BF on centrality and system size has been reported by several relativistic heavy-ion experiments [5, 6]. However, most of the current heavy-ion experiments are limited by the pseudorapidity range [5, 6, 7], it is impossible to quantitatively compare the results from the experiments with the coverage at different pseudorapidity ranges. The dependence of the BF on the pseudorapidity window is essential for understanding the physics of the BF [5, 6, 8, 9], and has been carefully studied by NA22 [9] and STAR experiments for hadron-hadron and relativistic heavy ion collisions, respectively.

The NA22 experiment has full 4π acceptance and excellent momentum resolution [9]. It is found in the experiment that the BF in π^+p and K$^+$p Collisions at 22 GeV is invariant under longitudinal boost over the whole rapidity range of produced particles, in spite of the non-boost-invariance of the single-particle density. Moreover, the BF of the whole rapidity range can be deduced from the BF properly scaled by the width of rapidity windows [9].

The STAR experiment covers a finite but relative wide pseudorapidity range. The scaling property of the BF in Au +Au collisions at 200 GeV is further observed in the experiment [10]. This scaling property of the balance function is also found in different p_{T} ranges of final state particles.

These results from both hadron-hadron and nuclear collisions indicate that charge balance of produced particles in strong interactions is boost-invariance in longitudinal phase-space, in contrary with the single particle density.

Therefore, it is interesting to see if those properties are taken into account in the models which are successfully described hadron-hadron and nuclear collisions, and how they associate with the mechanisms of particle production in the models.

II. CHARGE BALANCE FUNCTION AND IMPLEMENT MODELS

Charge balance function measures how the conserved electric charge compensate in the phase space, i.e., how the surrounding net charge are rearranged if the charge of a selected point changes[1]. In high energy collisions, the production of charged particles are constrained by charge balance in the phase space. The BF therefore provides a direct access to collision dynamics.

The BF has been originally defined in terms of a combination of four kinds of charge-related conditional densities in pseudorapidity [1]

$$\begin{aligned} B(\eta_1|\eta_2) \ = \ & \frac{1}{2}\left[\rho(+,\eta_1|-,\eta_2) - \rho(+,\eta_1|+,\eta_2)\right. \\ & \left. +\rho(-,\eta_1|+,\eta_2) - \rho(-,\eta_1|-,\eta_2)\right], \end{aligned} \quad (1)$$

where the notation $\rho(a,\eta_a|b,\eta_b)$ represents the ratio $\rho_{ab}(\eta_a,\eta_b)/\rho_b(\eta_b) = \langle n_{ab}(\eta_a,\eta_b)\rangle/\langle n_b(\eta_b)\rangle$ with a, b standing for + or − charged particles. Projecting to pseudorapidity difference $\delta\eta = \eta_1 - \eta_2$ in an pseudorapidity window η_w, it becomes [3, 5]

$$\begin{aligned} B(\delta\eta|\eta_{\mathrm{w}}) \ = \ & \frac{1}{2}\left[\frac{\langle n_{+-}(\delta\eta)\rangle - \langle n_{++}(\delta\eta)\rangle}{\langle n_+\rangle}\right. \\ & \left. + \frac{\langle n_{-+}(\delta\eta)\rangle - \langle n_{--}(\delta\eta)\rangle}{\langle n_-\rangle}\right] \end{aligned} \quad (2)$$

where $n_{ab}(\delta\eta)$ is the total number of pairs of opposite charged particles with pseudorapidity difference $\delta\eta$ in the pseudorapidity window $\eta_{\rm w}$. n_+ and n_- are the number of positively and negatively charged particles in the window $\eta_{\rm w}$, respectively. $\langle\cdots\rangle$ is the average over the whole event sample.

From the findings of the BF at NA22 [9] and STAR experiments [10], the BF is boost-invariant in the whole rapidity range in hadron-hadron collisions and may be in nuclear collisions as well. In the case, the properly scaled BF is corresponding to the BF of the whole pseudorapidity range and is deduced by

$$B_s(\delta\eta) = \frac{B(\delta\eta|\eta_{\rm w})}{1 - \frac{\delta\eta}{|\eta_{\rm w}|}} \qquad (3)$$

where $|\eta_{\rm w}|$ is the width of pseudorapidity window.

The PYTHIA 5.720 [11] is well set up for p+p collisions. It is a standard Monte Carlo generator with string fragmentation as hadronization scheme. Two versions of a multi-phase transport (AMPT) model [12] are used to study Au+Au collisions. One is the AMPT default and the other one is the AMPT with string melting. In both versions, the initial conditions are obtained from the HIJING model, and then the scattering among partons are given by ZPC. In the AMPT default model, the partons recombined with their parent strings when they stop interacting, and the resulting strings are converted to hadrons using the Lund string fragmentation model, while in the AMPT model with string melting, quark coalescence is used in combining partons into hadrons. The dynamics of the hadronic matter is described by ART model.

It is commonly believed that in relativistic heavy ion collisions, the charge ordering during the string fragmentation in elementary collisions is no longer valid, and it should be replaced by the quark-coalescence mechanism in hadronization [13]. So it is interesting to see whether the boost-invariance of the BF is sensitive to the mechanisms of hadronization.

In this paper, we firstly study the boost-invariance of the BF for p+p collisions at $\sqrt{s} = 22$ GeV and $\sqrt{s} = 200$ GeV using the PYTHIA, and for Au+Au collisions at $\sqrt{s} = 200$ GeV using two versions of the AMPT. The transverse momentum dependence of longitudinal scaling property of the BF is then examined in the models. The obtained results are compared with corresponding experimental data and discussed.

III. BOOST-INVARIANCE AND LONGITUDINAL SCALING OF THE BF

In order to demonstrate directly whether the BF is invariant under a longitudinal Lorentz transformation over the whole rapidity in hadron-hadron collisions, we choose four equal size ($|\eta_{\rm w}| = 3$) pseudorapidity windows locating at different positions $(-3, 0)$, $(-2, 1)$, $(-1, 2)$ and $(0, 3)$. The results for p+p collisions at $\sqrt{s} = 22$ GeV and $\sqrt{s} = 200$ GeV are shown in Fig. 1(a) and (b) respectively. The statistic errors are smaller than the markers. It is clear that the BF measured in four windows are approximately identical to each other at two incident energies. This indicates that the charge compensation is essentially the same in any longitudinally-Lorentz-transformed frame for p+p collisions in the PYTHIA model, consistent with the data from NA22 experiment. These results show that the string fragmentation mechanism implemented in PYTHIA well describes the production mechanisms of charge particles and their charge balance in longitudinal phase space.



FIG. 1: Upper panel: the $B(\delta\eta|\eta_{\rm w})$ in four pseudorapidity windows with equal size $|\eta_{\rm w}| = 3$ at the different positions for p+p collisions at (a) $\sqrt{s} = 22$ GeV and (b) $\sqrt{s} = 200$ GeV by PYTHIA model. Lower panel: the scaled balance function, $B_s(\delta\eta)$, deduced from the directly measured BF at six different sizes and positions of pseudorapidity windows for p+p collisions at (c) $\sqrt{s} = 22$ GeV and (d) $\sqrt{s} = 200$ GeV by PYTHIA model. The solid down triangle is the BF of the whole η range.

Fig. 1(c) and (d) are the scaled balance function $B_s(\delta\eta)$ at two incident energies. They are deduced from directly measured $B(\delta\eta|\eta_{\rm w})$ at six different pseudorapidity windows, $(-0.8, 0.8)$ (open circles), $(1, 3)$ (open triangles), $(-3, 1)$ (open squares), $(-2.4, 2.4)$ (open diamonds), $(0, 3)$ (open crosses), and $(-2, -1)$ (open stars). From the figures we can see that all the $B_s(\delta\eta)$ deduced from different windows are coincide with each other within errors, as expected from boost-invariance of the BF [4]. The solid down triangles in the same figures are the BF of the whole pseudorapidity range, $B(\delta\eta|\eta_\infty)$. It is close to the scaled balance function $B_s(\delta\eta)$. These results indicate that the scaled BF is in fact corresponding to the BF of the whole pseudorapidity range $B(\delta\eta|\infty)$ [4].

It is then interesting to see whether the boost-invariance of the BF is held in nucleus-nucleus collisions. STAR experiment only observe the boost-invariance of



FIG. 2: Upper panel: the $B(\delta\eta|\eta_{\rm w})$ in five pseudorapidity windows with equal size $|\eta_{\rm w}| = 2$ at the different positions for Au+Au collisions at $\sqrt{s} = 200$ GeV by (a) the AMPT default and (b) the AMPT with string melting. Lower panel: the scaled balance function, $B_s(\delta\eta)$, deduced from the directly measured BF at various pseudorapidity windows with different sizes and positions for Au+Au collisions at $\sqrt{s} = 200$ GeV by (c) the AMPT default and (d) the AMPT with sting melting.

BF in cental pseudorapidity range $-1 < \eta < 1$ [10], where the single particle distribution is almost flat, or boost-invariance. Now in model investigation, we can carefully examine the property in the whole pseudorapidity range.

The upper panel of Fig. 2 is the BF in five pseudorapidity windows with equal size $\eta_{\rm w} = 2$ at different positions $(-3, -1)$, $(-2, 0)$, $(-1, 1)$, $(0, 2)$ and $(1, 3)$. Where the Fig. 2(a) and (b) are the results from the AMPT default (v1.11) and the AMPT with string melting (v2.11), respectively. Both figures show that the BF is boost-invariance in pseudorapidity range (-3, 3) in two versions of the AMPT.

The lower panel of Fig. 2 is the scaled balance functions, which are obtained from directly measured BF at six different windows as indicated at legend of the figure, where the solid down triangles are the BF in pseudorapidity range (-4, 4). It shows that the scaled BF does not depend on the size and position of the windows, and corresponds to the BF of the whole pseudorapidity in two versions of the AMPT, consistent with the results of p+p collisions in the PYTHIA model.

IV. THE TRANSVERSE-MOMENTUM DEPENDENCE OF THE BOOST-INVARIANCE OF THE BF

The longitudinal property of boost invariance of BF comes from the special longitudinal interaction of charged particles under the constraint of global electric charge balance. Global electric charge conservation not only applies to all final-state charged particles, but also constrains particles which are produced at the same proper time of evolution. It is argued that the transverse-momentum of final-state particles may be roughly used as a scale of the proper time of their production in the expansion of nuclear collisions [14, 15, 16, 17]. Examining the $p_{\rm T}$ dependence of longitudinal property of the BF will provide direct access on whether particles in specified $p_{\rm T}$ range are consistent to be produced simultaneously with well balanced electric charge.

So we turn to check whether the boost-invariant of BF holds for particles in different $p_{\rm T}$ ranges. Fig. 3 shows the BF for p+p collisions at $\sqrt{s} = 22$ GeV and $\sqrt{s} = 200$ GeV from PYTHIA in three transverse momentum bins $(0 < p_{\rm T} < 0.2)$, $(0.2 < p_{\rm T} < 0.4)$, and $(p_{\rm T} > 0.2)$ GeV/c, respectively. These $p_{\rm T}$ bins are selected to make the multiplicity in each bin comparable. The result shows that the points at a given $\delta\eta$ in a restricted $p_{\rm T}$ interval are approximately coincide with each other, i.e., the boost-invariance of the BF hold in small $p_{\rm T}$ ranges. It indicates that particles produced at different $p_{\rm T}$ ranges are also boost-invariant for hadron-hadron collisions in the PYTHIA model.

The same study for Au+Au 200 GeV collisions from the two versions of the AMPT are presented in the upper and lower panels of Fig. 4, respectively. Where four $p_{\rm T}$ bins are, $(0.15, 0.4)$, $(0.4, 0.7)$, $(0.7, 1)$ and $(1, 2)$ GeV/c. From the upper panel of the figure, we can see that the BF of different pseudorapidity windows in each $p_{\rm T}$ bin are close to each other, in consistent with the data from STAR experiment [10]. However, in the AMPT with string melting, as shown in the lower panel of the figure, where the BF of different pseudorapidity windows are not as close to each other as those in the upper panel.

This is because in the AMPT with string melting, each parton in the evolution of nuclear collision has its own freeze-out time, which last a very long period after the interaction of two nucleus [18]. The particles in the same transverse-momentum range are not freezed-out simultaneously with well balanced charge, and therefore the longitudinal boost-invariance of the BF in small $p_{\rm T}$ ranges is violated. In the AMPT default, the partons recombined with their parent strings immediately after they stop interacting, and converted to hadrons. So the charge balance of the produced particles in the same $p_{\rm T}$ ranges is preserved and boost-invariance of the BF keeps.

V. SUMMARY

In the paper, we systematically study the longitudinal boost-invariance of charge balance function and its $p_{\rm T}$ dependence for p+p and Au+Au collisions using PYTHIA the AMPT models. It shows that charge balance function is boost-invariance in both hadron-hadron and nuclear interactions, in contrary to the single particle density. As



FIG. 3: For each of three $p_{\rm T}$ ranges, the $B(\delta\eta|\eta_{\rm w})$ in four pseudorapidity windows with equal size $|\eta_{\rm w}| = 3$ at the different positions for p+p collisions at $\sqrt{s} = 22$ GeV and $\sqrt{s} = 200$ GeV in upper and lower panels, respectively.



FIG. 4: For each of four $p_{\rm T}$ ranges, the $B(\delta\eta|\eta_{\rm w})$ in five pseudorapidity windows with equal size $|\eta_{\rm w}| = 2$ at the different positions for Au+Au collisions at $\sqrt{s} = 200$ GeV from the AMPT default (in upper panel) and the AMPT with string melting (in lower panel).

expected, this boost-invariance of the BF make the BF properly scaled by window size is independent of window and corresponds to the BF of the whole (pseudo)rapidity range. Therefore, the BF is a good measure free from the restriction of finite longitudinal acceptance.

It is further show that the boost invariance of the BF in specified $p_{\rm T}$ range is valid in PYTHIA for hadron-hadron collisions and the AMPT default for Au+Au collisions. While the AMPT with string melting fails to reproduce this property due to the different schemes at hadronization. So the $p_{\rm T}$ dependence of the longitudinal property of the BF may be served as a sensitive probe for charge balance in hadronization mechanism.

VI. ACKNOWLEDGMENTS

We thank Prof. Liu Lianshou and Dr. Yu Meiling for valuable discussions and remarks. This work is supported in part by the NSFC of China with project No. 10835005 and No. 10647124, and the MOE of China with project No. IRT0624 and No. B08033.

[1] D. Drijard *et al.*, Nucl. Phys. B **155** 269 (1979); B **166**, 233 (1980); I. V. Ajinenko *et al.*, *ibid.* C **43**, 37 (1989).

[2] R. Brandelik *et al.*, Phys. Lett. B **100**, 357 (1981); M. Althoff *et al.*, Z. Phys. C **17**, 5 (1983); H. Aihara *et al.*,

Phys. Rev. Lett. **53**, 2199 (1984); **57**, 3140 (1986) ; P. D. Acton *et al.*, Phys. Lett. B **305**, 415 (1993).

[3] S. A. Bass, P. Danielewicz and S. Pratt, Phys. Rev. Lett. **85**, 2689 (2000).

[4] S. Jeon and S. Pratt, Phys. Rev. C **65**, 044902 (2002).

[5] J. Adams *et al.* (STAR Collaboration), Phys. Rev. Lett. **90**, 172301 (2003); Gary. D. Westfall (for STAR Collaboration), J. Phys. G. **30**, S345 (2004).

[6] C. Alt *et al.* (NA49 Collaboration), Phys. Rev. C **71**, 034903 (2005); C. Alt *et al.* (NA49 Collaboration), arXiv: 0705.1122.

[7] K. Adcox *et al.* (PHENIX Collaboration), Phys. Rev. Lett. **89**, 082301 (2002).

[8] T. A. Trainor, hep-ph/0301122.

[9] M. R. Atayan *et al.* (NA22 Collaboration), Phys. Lett. B **637**, 39 (2006).

[10] Li Zhiming, Li Na, Liu Lianshou and Wu Yuanfang, Int. J Mod. Phys. E **16**, 3347 (2007).

[11] T. Sjöstrand, Comp. Phys. Commun. **82**, 74 (1994).

[12] Z. W. Lin, C. M. Ko, B. A. Li, B. Zhang and S. Pal, Phys. Rev. C **72**, 064901 (2005).

[13] A. Bialas, Phys. Lett. B **579**, 31 (2004); R. C. Hwa and C. B. Yang, Phys. Rev. C **70**, 024904 (2004).

[14] R. C. Hwa and Y. Wu, Phys. Rev. **C 60**, 054904 (1999).

[15] M. Asakawa, S. A. Bass, B. Müller and C. Nonaka, Phys. Rev. Lett. **101**, 122302 (2008).

[16] F. Grassi, Y. Hama, and T. Kodama, Phys. Lett. **B 355**, 9 (1995); Y. M. Sinyukov, S. V. Akkelin, and Y. Hama, Phys. Rev. Lett. **89**, 052301 (2002);

[17] M. Asakawa, S. A. Bass, B. Müller and C. Nonaka, Phys. Rev. Lett. **101**, 122302 (2008).

[18] Yu Meiling, Du Jiaxin, and Liu Lianshou, Phys. Rev. C (Rapid communication) **74**, 044906 (2006).